



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/31, C07K 14/255, G01N 33/50	A1	(11) International Publication Number: WO 98/03656 (43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/US97/12639 (22) International Filing Date: 18 July 1997 (18.07.97) (30) Priority Data: 60/022,191 19 July 1996 (19.07.96) US (71) Applicant (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; Morrill Hall, 100 Church Street, S.E., Minneapolis, MN 55455 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): RAJASHEKARA, Gireesh [IN/US]; Minn. Graduate Club, 2089 Carter Avenue, St. Paul, MN 55108 (US). NAGARAJA, Kakambi, V. [IN/US]; 2235 Milton Street North, Roseville, MN 55113 (US). KAPUR, Vivek [IN/US]; Apartment 108, 2572 Kenzie Terrace, St. Anthony, MN 55418 (US). (74) Agent: BRUESS, Steven, C.; Merchant, Gould, Smith, Edell, Welter & Schmidt, P.A., 3100 Norwest Center, 90 South Seventh Street, Minneapolis, MN 55402-4131 (US).		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: RECOMBINANT SEF14 FIMBRIAL PROTEIN FROM SALMONELLA (57) Abstract <p>A truncated SE fimbria antigen useful as an antigen for immunoassay diagnosis of <i>Salmonella enteritidis</i> (SE) infection or evidence of infection.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

RECOMBINANT SEF14 FIMBRIAL PROTEIN FROM SALMONELLA

Field of the Invention

The present invention relates to a method of
5 cloning and expressing a truncated form of a fimbrial gene
and the use of the truncated fimbrial gene product in an
immunodiagnostic assay and for immunoprophylaxis.

Background of the Invention

10 Foodborne infections cause an estimated 6.5
million cases of human illness and 9000 deaths annually in
the United States alone. Bacterial infections by
Salmonella are the most commonly reported cause of
foodborne outbreaks. *Salmonella enteritidis* (SE) is the
15 dominant *Salmonella* serotype isolated from cases of food
poisoning. Many of these outbreaks are thought to be due
to infected poultry products, particularly eggs and egg
products.

The best way to prevent infection in human
20 populations is to diagnose and treat the infected animal
prior to human consumption. Because the greatest threat of
food poisoning from *Salmonella* is from poultry products,
there is a need for a method to detect birds that are
infected with SE.

25 Some current diagnostic methods rely on
conventional bacteriologic cultures. However, these
procedures are relatively slow, often taking up to 3 to 4
days to provide even a presumptive diagnosis.
Additionally, the great susceptibility of SE to physical
30 and chemical factors such as desiccation, radiation, low
temperature, heating, or chemical preservatives, causes
traditional bacteriologic culture methods to generally have
a low sensitivity. Consequently, many birds or animals

that are infected with SE are often not detected when conventional bacterial cultures are used.

Other diagnostic methods rely on the detection of serum antibodies specific to SE. Although several serological methods such as micro-agglutination, serum plate agglutination, latex particle agglutination, microantiglobulin, ELISA have previously been employed, these assays lack either the sensitivity or specificity necessary to detect SE infected birds, or the tests are too difficult to perform in a routine laboratory or field setting. Consequently, widespread application of these tests for the detection of SE infections has been impractical.

A useful antigenic determinant that is found on many species of *Enterobacteriaceae* are fimbriae, proteinaceous filamentous surface structures composed of protein subunits called fimbrin. Upon infection, birds make antibodies to this SE fimbrial antigen. Therefore, the SE fimbrial antigen is useful in a diagnostic assay for the presence of SE in poultry.

SE is known to have at least four distinct fimbria, designated Sef14, Sef17, Sef18 and Sef21. These proteins are encoded by *SefA*, *AgfA*, *SefD* and *FinA* genes, respectively.

Although the gene encoding Sef14 has been identified and its DNA nucleotide sequence determined (Trucotte and Woodward, *Journal of General Microbiology*, 139:1477-1485 (1993)), an effective diagnostic method using this surface antigen has not been developed, partially due to the difficulty of efficiently producing the fimbriae proteins in purified form and in large quantities. Additionally, expression of Sef14 fimbriae by cultured

Salmonella enteritidis is highly dependent on the growth medium composition. In a study by Thorns et al., *International Journal of Food Microbiology*, 21:47-53 (1994), only peptone water pH 7.2 supported the expression of Sef14 by all *Salmonella enteritidis* strains examined. Consequently, previous diagnostic assays using Sef14 have used antibodies against Sef14 and not the antigen itself.

Hence, there is a need for a sensitive, specific and routine antigen and method to reliably detect SE infection in birds, preferably a method that is easily adaptable to large-scale screening of poultry flocks.

Summary of the Invention

The present invention provides a sensitive, specific, routine antigen and assay to reliably detect SE-infected animals. Specifically, the present invention provides a truncated form of the Sef14 antigen that can be easily produced in purified form and in large quantities and used in the method of the invention. The novel Sef14 antigen, when coupled to a substrate such as latex beads, provides a diagnostic assay for SE, particularly useful in large-scale screening of poultry flocks.

Brief Description of the Figures

Figure 1 is a photograph showing a SDS-PAGE of the recombinant Sef14 (rSef14) fragment (arrow).

Figure 2 is a photograph showing a Western blot of the rSef14 fragment probed with anti-Sef14 antibody (lane 1) and anti-tag (T7) antibody (lane 2).

Figure 3 is a photograph showing results of a rSef14-latex agglutination assay for SE infection in

chickens exposed to *S. enteritidis* (A), *S. pullorum* (B), and serum-free antigen control (C).

Figure 4 is a photograph showing results of a rSef14-latex agglutination assay for SE infection in

5 chickens exposed to *S. enteritidis* (A), *S. gallinarum* (B), *S. pullorum* (C), *S. typhimurium* (D), *C. arizonae* (E), *E. coli* (F), serum free antigen control (G), and serum control (H).

Figure 5 is a graph showing the percentage of
10 chickens testing positive for anti-SE antibodies during 4 weeks post-inneculation. The five bars at each week represent inneculation with 10^4 , 10^6 , 10^8 , 10^{10} , and control (no cells).

Figure 6 is a graph showing the antibody titres
15 of chicken sera samples testing positive for anti-SE antibodies.

Figure 7 is a graph showing the antibody titres
of chicken egg yolk samples testing positive for anti-SE antibodies.

Detailed Description of the Invention

The present invention is directed to a method for diagnosing *Salmonella enteritidis* infection or evidence of infection in an animal, particularly poultry, using a
25 recombinant truncated fimbrial antigen.

"Infection" means active colonization of the animal by SE organisms. "Evidence of infection" means a prior history of colonization by SE in the animal, although active colonization is not present. Diagnosis of active
30 infection is needed to protect against contamination of food supplies, whereas diagnosis of prior infection is

needed to alert against new infection or to trace the source of infection in a flock.

Fimbrial Proteins

5 Fimbriae are proteinaceous filamentous surface structures composed of protein subunits called fimbrin. These proteinaceous structures are thought to be virulence factors which mediate specific attachment to host cell mucosal surfaces. They are present in most enteric
10 bacteria capable of invading host cells.

Salmonella enteritidis has four distinct fimbriae: Sef14, Sef17, Sef18 and Sef21 which are encoded by *sefA*, *agfA*, *sefD* and *fimA* genes, respectively. Sef14 is unique with only limited distribution in the genus. In
15 contrast, all other fimbrial proteins are widely distributed in the genus. Thus, they have limited use as diagnostic reagents for SE detection.

Cloning and Expression of Sef14

20 In the present invention, a truncated form of the Sef14 antigen retaining the antigenic character of the entire protein has been produced. Unlike the complete protein, however, the truncated form can be easily produced in purified form and in large quantities, without special
25 growth medium requirements.

PCR technology is used to produce the truncated Sef14 protein by amplification with suitable primers.

Primers are selected to amplify the gene encoding Sef14 in a region downstream of the encoded signal peptide, e.g., downstream of about nucleotide 145 of the DraI
30 genomic fragment shown in Figure 1 of Turcotte and Woodward, *Supra*. Preferably, the PCR primers include

additional nucleotides at the 5' ends, encoding specific restriction enzyme recognition sequences, for ease of purification. For example, useful primers for amplifying that portion of the *sefA* gene encoding an immunogenic Sef14 fragment downstream of the signal peptide are shown below:

GGGAATTGCTGGCTTTGTTGGTAACA	SEQ ID NO:1
GGGCTCGAGTTAGTTTGTGATACTGAACGTA	SEQ ID NO:2

After a truncated gene sequence encoding Sef14 is produced, it can be cloned into a host using a plasmid or phage as a vector. Typically, the expression of Sef14 fimbriae by cultured *Salmonella enteritidis* is highly dependent on the growth medium composition (Thorns et al, *International Journal of Food Microbiology*, 21:47-53 (1994)), and it is typically difficult to produce large quantities. However, a truncated form of Sef14 having at least the signal peptide removed is expressed in host systems such as *E. coli* without these difficulties.

Truncated Sef14 Antigen

Because the truncated Sef14 protein retains the antigenic characteristics of the complete protein, it is useful in various immunological methods. For example, the inventive antigen is useful in antibody binding immunoassays such as assays to detect the presence of antibodies against SE in a sample. Suitable binding assays include ELISA, wherein the recombinant Sef14 antigen is bound to a surface and exposed to antibodies against SE. To detect the presence of bound anti-SE antibodies, a marker such as an enzyme-linked secondary antibody is then added.

An agglutination assay using truncated Sef14 antigen-coated latex beads is preferred. In the agglutination reaction, antigen-coated latex beads form detectable clusters when exposed to antibodies against SE. This preferred assay is described more fully in Example 4, below.

Diagnostic Assays

The assays described above can be used to detect the presence of antibodies to *Salmonella enteritidis*. Preferably, the assays are used to determine whether or not an animal, e.g. a poultry animal such as a chicken or turkey, is infected with SE. Animal fluid such as blood or serum can be used in a diagnostic assay. If an animal is infected with SE, the animal will typically produce anti-SE antibodies. The recombinant Sef14 antigen is used to detect the presence of anti-SE antibodies, SE infection or the SE organism itself. Diagnostic assays such as these are particularly useful in birds. More particularly, diagnostic assays are useful in detecting SE infections in chicken or turkey to prevent foodborne illness by poultry consumption.

Vaccine

Passive immunization with anti-Sef14 antibodies has been shown to reduce *Salmonella enteritidis* colonization (Peralta et al. 1994). Additionally, Sef14 can induce a T-cell immune response (Ogunniyi et al 1994). Because the truncated Sef14 antigen exhibits these immunological activities, can be produced in large quantities, and does not have the cumbersome growth requirements of the complete protein, the truncated Sef14

antigen is also useful as a vaccine to confer immunity against SE. Preferably, the truncated Sef14 antigen is used as a vaccine in poultry to prevent foodborne illnesses.

5

EXAMPLES

The invention may be better understood with reference to the following examples which are not intended to limit the invention.

10

Example 1

Isolation of *S. enteritidis* genomic DNA

15

S. enteritidis was grown overnight at 37°C in Luria-Bertani (LB) broth. Genomic DNA was extracted as described (Sambrook, et al., 1989) using standard methods with minor modifications. In brief, bacterial cells were pelleted by centrifugation at 13,000 x g for 3 minutes, washed/suspended in 1 ml of 1 M NaCl, centrifuged for 5 minutes at 13,000 x g, and the pellet resuspended in 1 ml TE buffer (50 mM Tris-HCl, 50 mM EDTA, pH 7.8). The sample was next incubated with 5 µl of lysozyme (50 mg/ml) (Sigma Chemical Co., St. Louis, MO) and 0.3 mg/ml RNase A (Sigma) at 37°C for 30 minutes. To this suspension, 1% sarkosyl and 0.6 mg/ml of proteinase K (Sigma) were added, and the mixture incubated at 37° for 1 hour. Following incubation, chromosomal DNA was extracted twice with an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1) and once with chloroform:isoamyl alcohol (24:1). Genomic DNA in the aqueous phase was precipitated at -20°C with two volumes of absolute ethanol and 0.1 volume of 3 M sodium acetate, and

20

25

30

pelleted by centrifugation at 13,000 x g for 5 minutes. The pellet was then washed twice with 70% ethanol, air dried, and suspended in TE Buffer (10mM Tris-HCl and 1mM EDTA pH 8.0). Total DNA was quantitated
5 spectrophotometrically at A₂₆₀ nm.

Example 2

Cloning of *sefA* gene fragment

Oligonucleotide primer selection and synthesis:

10 Oligonucleotide primers corresponding to an internal fragment (64-498 bp) of the open-reading frame of the *sefA* gene were used for PCR amplification. Additional bases were added to the 5' end of each primer in order to confer a recognition sequence for either *EcoRI* (forward
15 primer) or *XhoI* (reverse primer). The oligonucleotide primers were obtained from Integrated DNA technologies Inc., Ames, IA. The DNA sequences for the forward and reverse primers are shown below:

<u>GGGAATTC</u> GCTGGCTTTGTTGGTAACA	SEQ ID NO:1
<u>GGGCTCGAGT</u> TAGTTTTGATACTGAACGTA	SEQ ID NO:2

20 Additional nucleotides added to the 5' end of the primers are underlined.

PCR amplification of *sefA* gene fragment.

Amplification reactions were performed in 30 µl volumes with 30 pmol of each primer and 5 mM MgCl₂. The
25 reagents and enzymes used for PCR were obtained either from Boehringer Mannheim (Indianapolis, IN) or Perkin Elmer (Foster City, CA). One hundred ng of genomic DNA was used as a template for PCR amplification with the following parameters: an initial denaturation at 94°C for 5 minutes,
30 followed by 35 cycles of denaturation (94°C for 1.5

minutes), annealing 52°C for 1 minute) and extension (72°C for 2 minutes), and a final extension of 15 minutes at 72°C. All amplification reactions were performed in a Perkin-Elmer Cetus DNA thermal cycler (Model 480). The PCR products were analyzed on a 1% agarose gel, stained with ethidium bromide (0.5µg/ml), and photographed under UV light.

PCR products were gel extracted (Qiagen Inc., Chatsworth, CA), quantitated spectrophotometrically, at 260 nm, and cloned directly into pGEM-T vector (Promega, Madison, WI). Following ligation, 2µl of the reaction products were transformed into *E. coli* DH5α cells (Gibco BRL, Gaithersburg, MD) by the heat shock method. Recombinant colonies were selected on ampicillin/IPTG-Xgal containing plates and screened for the presence of the appropriate insert by restriction analysis.

Nucleotide sequence analysis

A bacterial colony containing the recombinant plasmid with the *rSefA* fragment was grown in LB-ampicillin media, and the plasmid extracted using Qiagen plasmid extraction kit (Qiagen). The nucleotide sequence of the insert was determined using oligonucleotide primers specific to the vector sequence by automated DNA sequencing at the University of Minnesota Advanced Genetic Analysis Center. The insert was sequenced in its entirety in both orientations, and the amino acid sequence deduced using the standard genetic code (DNA*, Madison, WI). Sequencing results are shown below for nucleotide and deduced amino acid sequences of the insert (Seq.ID.NO:5), together with a tag sequence added during the subcloning of the fragment into the pET/abc expression vector (Seq.ID.NO:3). The

added tag sequence at the 5' end, provides a Histidine-rich portion to facilitate purification of the sequence on nickel columns, as well as an antigenic region that specifically binds the T7 anti-tag antibody provided with the pET/abc vector kit.

Nucleic Acid Sequence encoding rSefA fragment

SEQ ID NO:3

```

ATG GGC AGC AGC CAT CAT CAT CAT CAT CAC AGC AGC GGC CTG GTG 45
CCG CGC GGC AGC CAT ATG GCT AGC ATG ACT GGT GGA CAG CAA ATG 90
GGT CGC GGA TGG GAA TTC GCT GGC TTT GTT GGT AAC AAA GCA GTG 135
GTT CAG GCA GCG GTT ACT ATT GCA GCT CAG AAT ACA ACA TCA GCC 180
AAC TGG AGT CAG GAT CCT GGC TTT ACA GGG CCT GCT GTT GCT GCT 225
GGT CAG AAA GTT GGT ACT CTC AGC ATT ACT GCT ACT GGT CCA CAT 270
AAC TCA GTA TCT ATT GCA GGT AAA GGG GCT TCG GTA TCT GGT GGT 315
GTA GCC ACT GTC CCG TTC GTT GAT GGA CAA GGA CAG CCT GTT TTC 360
CGT GGG CGT ATT CAG GGA GCC AAT ATT AAT GAC CAA GCA AAT ACT 405
GGA ATT GAC GGG CTT GCA GGT TGG CGA GTT GCC AGC TCT CAA GAA 450
ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT AAA TCG ACC CTG CCA 495
GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG CAG TAT CAA AAC 540
TAA CTC GAG CCC 552

```

*Additional amino acid residues added to the amino terminus to facilitate protein purification and cloning are underlined.

Deduced amino acid sequence of rSefA protein fragment*

Seq. ID NO: 4

```

MGSSHHHHHHSSGLVPRGSHMASMTGGQOMGRGSEFAGFVGNGKAVVQAAVT
IAAQNTTSANWSQDPGFTGPAVAAGQKVGTL SITATGPHNSVSIAGKGASVSGG
VATVPFVDGQGQPVFRGRIQGANINDQANTGIDGLAGWRVASSQETLNPVTT
FGKSTLPAGTFTATFYVQQYQN

```

*Additional amino acid residues added to the amino terminus to facilitate protein purification and cloning are underlined.

It is understood that the amino acids added to the N-terminus of the Spf14 antigen are optional, and used for ease of cloning and purification. The amino acid sequence in the absence of these added residues (Sequence ID No:6)

with or without other added residues for cloning or purification procedures, for example, are similarly useful as antigens in the diagnostic assays of the invention.

5 **Subcloning *sefA* gene fragment into an expression vector**

 The pGEM-T plasmid carrying *sefA* fragment was double digested with *EcoRI* and *XhoI*, and the digested products gel purified (Qiagen) and cloned into *EcoRI* and *XhoI* digested pET/abc expression vectors (Novagen Inc.,
10 Madison, WI). Ligation products (2 μ l) from each of the reactions were transformed into *E. coli* BL21(DE3)pLyS cells by heat shock method. Recombinant clones were cultured on kanamycin and chloramphenicol containing plates, and analyzed by restriction enzyme analysis.

15

rSefA fragment expression

 The recombinant clones were selected based on restriction enzyme analysis with *EcoRI* and *XhoI* digestion, selecting those clones yielding appropriately sized
20 fragments as compared with a vector control. Selected clones were analyzed for rSefA fragment expression. Briefly, a single colony from each (pETabc/*SefA* fragment) freshly streaked plate was picked and inoculated to 50 ml LB broth containing appropriate antibiotics and incubated
25 with shaking at 200 rpm at 37°C until the OD₆₀₀ reached 0.6. Cultures were induced with IPTG (0.4 mM) and incubated for an additional 3 hours. Following incubation, the cells were pelleted and resuspended in 5 ml of TE buffer (50mM Tris-HCl pH 8.0, 2mM EDTA) and incubated with 25 μ l of
30 lysozyme (50 mg/ml) and 100 μ l of 1% Triton X-100 for 20 minutes at 30°C. The samples were sonicated until they were no longer viscous, and centrifuged at 39,000 x g for

20 minutes. The supernatant was passed through a 0.45 μ m membrane filter, and stored at -20°C until further use.

SDS-PAGE analysis

5 The cell lysates were next analyzed by SDS-PAGE for the presence of the rSefA fragment by mixing with an equal volume of 2x SDS solubilization buffer separating on 12% polyacrylamide gels, and staining with Coomassie blue. The results are shown in lane 1 of Figure 1 which contains
10 the total protein produced by the vector and contained in the cell lysates.

Western blot analysis

 The lysates were separated on 12% polyacrylamide
15 gels and transferred onto a nitrocellulose membrane using Transblot apparatus (Bio-Rad laboratories, Hercules, CA). Following transfer, the membrane was blocked with 3% BSA in phosphate buffered saline (PBS) and stained with either T7 anti-tag antibody (Novagen) or rabbit anti-Sef14 specific
20 antibody (kindly provided by Dr. W. W. Kay, University of Victoria, BC, Canada). The membrane was washed and stained with anti-rabbit IgG/HRP conjugate and treated with developing reagent (Amersham lif sciences, Inc., USA) for 1
25 minute, exposed to X-ray film, and the radiograph developed. The results are shown in Figure 2, where lane 1 is probed with anti-Sef14 antibody, and lane 2 with T7 anti-tag antibody.

30 Purification of rSef14 fragment protein by column chromatography and electroelution

 The recombinant Sef14 protein fragment produced in the cell lysates described above was purified by binding

of the Histidine-rich tag to nickel columns as described by the manufacturer (Novagen). Briefly, the cells were induced and extract was prepared as described above except that the induced cells were suspended in Tris buffer without EDTA. The cell lysate was passed through nickel columns and washed sequentially with binding buffer (5 mM imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9) and wash buffer (60 mM imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9). The bound protein was eluted using elution buffer (1 M imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9), quantitated using a Bio-Rad protein assay kit (Bio-Rad laboratories, Hercules, CA), and analyzed by SDS-PAGE. (See Figure 1, lane 2, where the arrow indicates the rSef14 fragment at about 19 KDa.) Since the column purified recombinant material contained traces of non-specific proteins, the appropriate rSef14 fragment was further purified by cutting the rSef14 fragment from the gel and electroelution (Bio-Rad) following the manufacturer's suggested protocol. The electroeluted fragment is shown in lane 4 of Figure 1 (at arrow).

Example 3

Covalent coupling of rSef14 to blue-dyed latex beads

The electroeluted rSef14 protein fragment was coupled to either 0.5 μ m or 1.0 μ m blue-dyed latex beads (Polysciences Inc., Warrington, PA) by glutaraldehyde method. Briefly, 1 ml of 2.5% suspension of the beads were washed with PBS (pH 7.4), pelleted by centrifugation and resuspended in 1 ml of 8% glutaraldehyde (EM grade) in PBS, and incubated overnight with gentle end-to-end mixing at room temperature. Following glutaraldehyde treatment, the beads were pelleted, washed with PBS three times and

incubated with 500 µg of purified rSef14 fragment for 5 hours at room temperature with gentle end-to-end mixing. The beads were pelleted, and incubated with 1 ml of 0.5 M ethanolamine in PBS for 30 minutes at room temperature with gentle end-to-end mixing. The mixture was then treated with 1 ml of 10 mg/ml BSA in PBS for 30 minutes at room temperature, centrifuged and the pellet resuspended in 1 ml PBS (pH 7.4), containing 10 mg/ml BSA, 0.1% NaN₃, and 5% glycerol, and stored at 4°C to form rSef14 - fragment coated latex beads for use in agglutination assays.

Example 4

rSef14-latex bead agglutination test

Bacteria was administered to chickens by either injection, intratracheal or oral administration of 10⁷ colony forming units (CFU) of either *S. enteritidis*, *S. pullorum*, *S. arizonae*, *S. typhimurium*, *S. gallinarum*, or *E. coli*. After about two to three weeks exposure, serum was collected and used to evaluate the sensitivity and specificity of the rSef14-latex beads in an agglutination assay for anti-SE antibody binding. A total volume of 7.5 µl of rSef14 fragment coated latex beads, produced as described for Example 3, were mixed with an equal volume of chicken serum collected from birds exposed to various pathogens, as described above. The presence of agglutination, visually seen as a loss of intense blue color in the sample (i.e., lightening of color as the coated beads agglutinate or form a lattice). Absence of the agglutination reaction was visualized by the remaining intense blue color of the dyed beads in a homogeneous suspension. Positive or negative agglutination reaction

was recorded after two minutes. The results are shown in Figures 3 and 4.

In figure 3, intense blue color (negative result) is seen in test samples B and C (*S. pullorum* and the serum-free antigen control). In contrast, a positive agglutination result is seen in test Sample A, (*S. enteritidis*), as a pale blue, diffuse agglutination pattern.

In figure 4, a positive agglutination reaction is seen in sample A (*S. enteritidis*) and in sample H (serum control). No agglutination reaction is seen in the samples B-G containing serum animals exposed to the following pathogens: *S. gallinarium* (B), *S. pullorum* (C), *S. typhimurium* (D), *S. arizonae* (E), *E. coli* (F), and serum free antigen control (H).

Example 5

Detection of anti-S.E. antibodies in infected chickens

To confirm the specificity of the assay of the invention, forty SPF chickens (age 4weeks) were inoculated with various species of *Salmonella*. A suspension of 10^9 CFU in PBS was administered by injection. A booster dose of 10^9 CFU was administered orally two weeks later. Serum samples were taken at weekly intervals and assayed for the presence of anti-SE antibodies.

Serum samples were assayed by the standard Serum Plate Test (SPT) as described in Schaffer and MacDonald, 1931, _____, ____:236-240. Slides were visually scored (+ or -) for the presence of agglutination, indicating antigen-antibody binding. Samples were also assayed by the Microtiter Plate Test (MPT) as described in Williams and Whittemore, 1971 *Applied Microbiology* 21:394-399. These

two assays are standard screening methods for the detection of Salmonella, using *S.pullorium* as a whole-cell antigen, and are not specific for SE, as shown in the table below.

To demonstrate the specificity of the assays of the invention, serum samples were assayed using the latex agglutination test (LAT) described above for Example 4, which utilized the truncated Spf14 antigen coupled to latex beads. Serum samples were also assayed for anti-SE antibodies by ELISA. In the ELISA, the truncated Spf14 antigen prepared as described for Example 3, was coated onto polystyrene plates. Antigen-coated plates were exposed to serum samples to permit binding of anti-SE antibodies to the antigen. The bound antigen-antibody complexes were washed, and then incubated with anti-chicken antibody coupled to biotin. The complex was then exposed to strep-avidin for signal detection.

Results are shown in the table below. The LAT and Elisa assays demonstrated a useful specificity for the detection of SE. Of the organisms tested, only *S.dublin*, a bovine pathogen, demonstrated cross-reactivity in the assays.

Species	SPT	MT	LAT	ELISA
<i>S. enteritidis</i>	+	+	+	+
<i>S. gallinarum</i>	+	+	-	-
<i>S. pullorum</i>	+	+	-	-
<i>S. dublin</i>	+	+	+	+
<i>S. berta</i>	+	+	-	-
<i>S. typhimurium</i>	-	+	-	-
<i>E. coli</i>	-	-	-	-
Control (no cells)	-	-	-	-

Example 6**Specificity of anti-SE assay**

The ELISA assay for detecting anti-SE antibodies described above for Example 6 was tested for specificity using a panel of antisera against known pathogenic organisms. Each sera was assayed in the anti-SE ELISA. No crossreactivity was observed with any of the tested antisera.

10

Antisera	ELISA	Antisera	ELISA
Pox	-	MG	-
Reo	-	NDV	-
Rev	-	CAV	-
SB-1	-	HVT	-
IBDV	-	IBV	-
ILT	-	<i>S.typhimurum</i>	-
LLA	-	<i>S.gallinarum</i>	-
LLB	-	<i>S.pullorum</i>	-
MS	-		

15

Example 7**Sensitivity of ELISA for detection of SE**

Fifty white leghorn layer chickens (5 weeks old) were orally inoculated in a single exposure with varied amounts of SE, from 10^4 to 10^{10} CFU in PBS. Serum samples were collected at weekly intervals for up to seven weeks. Eggs were collected for egg yolk antibody detection.

20

Samples were analyzed for detection of anti-SE

antibodies using the ELISA described above for Example 6. As shown in Figure 5, control chickens showed no positive reaction in the ELISA assay. Approximately 40-80% of chickens exposed to 10^4 , 10^6 , 10^8 , and 10^{10} CFU of SE tested positive for anti-SE antibodies during the first four weeks post-inneculation. From 4-7 weeks post-inneculation, the data stabilized at about 45% positive detection of anti-SE antibodies.

Antibody titers in the sera and egg yolks of chickens exposed to 10^4 , 10^6 , and 10^8 CFU of SE and testing positive in the ELISA for anti-SE antibodies are shown in Figures 6 and 7.

These data demonstrate specific detection of anti-SE antibodies using recombinant Sef14-antigen coated latex beads in an agglutination assay and using the antigen as a capture agent in an ELISA. These assays provide a sensitive and specific diagnostic tool for the detection of anti-SE antibodies in animals and for the diagnosis of SE infection.

20

SEQUENCE LISTING

- (1) GENERAL INFORMATION
- (i) APPLICANT: REGENTS OF THE UNIVERSITY OF MINNESOTA
- (ii) TITLE OF THE INVENTION:
RECOMBINANT FIMBRIAL PROTEIN
- (iii) NUMBER OF SEQUENCES: 6
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Merchant, Gould, Smith, Edell, Welter & Schmidt
(B) STREET: 3100 Norwest Center, 90 South Seventh St
(C) CITY: Minneapolis
(D) STATE: MN
(E) COUNTRY: USA
(F) ZIP: 55402
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ Version 2.0
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE: 18-JUL-1997
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 60/022,191
(B) FILING DATE: 19-JUL-1996
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Kettelberger, Denise M
(B) REGISTRATION NUMBER: 33,924
(C) REFERENCE/DOCKET NUMBER: 600.335WO01
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 612/371-5268
(B) TELEFAX: 612/332-9081
(C) TELEX:

22

GGA TGG GAA TTC GCT GGC TTT GTT GGT AAC AAA GCA GTG GTT CAG GCA	144
Gly Trp Glu Phe Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala	
35 40 45	
GCG GTT ACT ATT GCA GCT CAG AAT ACA ACA TCA GCC AAC TGG AGT CAG	192
Ala Val Thr Ile Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln	
50 55 60	
GAT CCT GGC TTT ACA GGG CCT GCT GTT GCT GCT GGT CAG AAA GTT GGT	240
Asp Pro Gly Phe Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly	
65 70 75 80	
ACT CTC AGC ATT ACT GCT ACT GGT CCA CAT AAC TCA GTA TCT ATT GCA	288
Thr Leu Ser Ile Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala	
85 90 95	
GGT AAA GGG GCT TCG GTA TCT GGT GGT GTA GCC ACT GTC CCG TTC GTT	336
Gly Lys Gly Ala Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val	
100 105 110	
GAT GGA CAA GGA CAG CCT GTT TTC CGT GGG CGT ATT CAG GGA GCC AAT	384
Asp Gly Gln Gly Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn	
115 120 125	
ATT AAT GAC CAA GCA AAT ACT GGA ATT GAC GGG CTT GCA GGT TGG CGA	432
Ile Asn Asp Gln Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg	
130 135 140	
GTT GCC AGC TCT CAA GAA ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT	480
Val Ala Ser Ser Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly	
145 150 155 160	
AAA TCG ACC CTG CCA GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG	528
Lys Ser Thr Leu Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln	
165 170 175	
CAG TAT CAA AAC TAACTCGAGC CC	552
Gln Tyr Gln Asn	
180	

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 180 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

23

```

Met Gly Ser Ser His His His His His His Ser Ser Gly Leu Val Pro
 1           5           10           15
Arg Gly Ser His Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg
          20           25           30
Gly Trp Glu Phe Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala
          35           40           45
Ala Val Thr Ile Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln
          50           55           60
Asp Pro Gly Phe Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly
65           70           75           80
Thr Leu Ser Ile Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala
          85           90           95
Gly Lys Gly Ala Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val
          100          105          110
Asp Gly Gln Gly Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn
          115          120          125
Ile Asn Asp Gln Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg
          130          135          140
Val Ala Ser Ser Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly
145          150          155          160
Lys Ser Thr Leu Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln
          165          170          175
Gln Tyr Gln Asn
          180

```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 435 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 1...432
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

GCT GGC TTT GTT GGT AAC AAA GCA GTG GTT CAG GCA GCG GTT ACT ATT      48
Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala Ala Val Thr Ile
 1           5           10           15

GCA GCT CAG AAT ACA ACA TCA GCC AAC TGG AGT CAG GAT CCT GGC TTT      96
Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln Asp Pro Gly Phe
          20           25           30

ACA GGG CCT GCT GTT GCT GCT GGT CAG AAA GTT GGT ACT CTC AGC ATT     144
Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly Thr Leu Ser Ile
          35           40           45

```

24

ACT GCT ACT GGT CCA CAT AAC TCA GTA TCT ATT GCA GGT AAA GGG GCT	192
Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala Gly Lys Gly Ala	
50 55 60	
TCG GTA TCT GGT GGT GTA GCC ACT GTC CCG TTC GTT GAT GGA CAA GGA	240
Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val Asp Gly Gln Gly	
65 70 75 80	
CAG CCT GTT TTC CGT GGG CGT ATT CAG GGA GCC AAT ATT AAT GAC CAA	288
Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn Ile Asn Asp Gln	
85 90 95	
GCA AAT ACT GGA ATT GAC GGG CTT GCA GGT TGG CGA GTT GCC AGC TCT	336
Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg Val Ala Ser Ser	
100 105 110	
CAA GAA ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT AAA TCG ACC CTG	384
Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly Lys Ser Thr Leu	
115 120 125	
CCA GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG CAG TAT CAA AAC T	433
Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln Gln Tyr Gln Asn	
130 135 140	
AA	435

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala Ala Val Thr Ile	
1 5 10 15	
Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln Asp Pro Gly Phe	
20 25 30	
Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly Thr Leu Ser Ile	
35 40 45	
Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala Gly Lys Gly Ala	
50 55 60	
Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val Asp Gly Gln Gly	
65 70 75 80	
Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn Ile Asn Asp Gln	
85 90 95	
Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg Val Ala Ser Ser	
100 105 110	
Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly Lys Ser Thr Leu	
115 120 125	
Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln Gln Tyr Gln Asn	
130 135 140	

WE CLAIM:

1. A method for detecting anti- *Salmonella enteritidis* antibodies in animals, the method comprising:

5 reacting a sample obtained from an animal with a truncated Sef14 antigen under conditions to permit anti-Sef14 antibodies to bind the antigen, the truncated antigen having at least the native Sef14 signal peptide removed; and

10 correlating antibody-antigen binding with the presence of anti-SE antibodies in the sample.

2. A method for diagnosing *Salmonella enteritidis* infection in animals, the method comprising:

15 reacting a sample obtained from an animal with a truncated Sef14 antigen under conditions to permit anti-Sef14 antibodies to bind the antigen, the truncated antigen having at least the native Sef14 signal peptide removed; and

20 correlating antibody-antigen binding with *Salmonella enteritidis* infection.

3. The method of claim 1, wherein the truncated Sef14 antigen has the amino acid sequence of Sequence ID. No. 4.

25

4. The method of claim 1, wherein the truncated Sef14 antigen has the amino acid sequence of Sequence ID. No.6.

5. The method of claim 1, wherein said antigen is fixed to
30 an inert surface prior to said reacting.

6. A Sef14 antigen consisting essentially of the amino acid sequence of Sequence I.D. No. 6.


5 7. An assay kit for the detection of anti-*Salmonella enteritidis* antibodies comprising an Sef14 antigen consisting essentially of the amino acid sequence of Sequence ID No. 6.

10 8. The assay kit of claim 7, wherein the antigen consists essentially of the amino acid sequence of Sequence ID. No. 4.

15 9. An antigen for stimulating the production of anti-*Salmonella enteritidis* antibodies comprising the amino acid sequence of Sequence ID No. 4 or 6.

20 10. The method of detecting anti-*Salmonella enteritidis* antibodies described in any of the foregoing claims, wherein the animal samples are obtained from fowl, and particularly from chickens or turkeys.

Figure 1

SDS-PAGE of rSef14 fragment protein

1. BL21DE3 (pET/sefA) before purification
2. Purified rSef14 fragment protein
3. LMW marker
4. rSef14 after electroelution

Figure 2

**Western blot -- rSef14 fragment probed
with Sef14 monospecific polyclonal
antibody**




Figure 3

Latex agglutination test




C

B

A

Figure 4


1. Latex agglutination test.



5/7

Figure 5


Sensitivity of rSEF14-LAT Chicken serum



6/7

Figure 6


Sensitivity of SEF-14 ELISA Using Chicken Sera



7/7

Figure 7

Sensitivity of SEF-14 ELISA Using Yolk



INTERNATIONAL SEARCH REPORT

Inte .onal Application No

PCT/US 97/12639

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/31 C07K14/255 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THORNS CJ ET AL: "Development and application of enzyme-linked immunosorbent assay for specific detection of Salmonella enteritidis infections in chickens based on antibodies to SEF14 fimbrial antigen." J CLIN MICROBIOL, APR 1996, 34 (4) P792-7, UNITED STATES, XP002047275 see page 792 - page 793; table 1 ---	1,2,5,6, 10
X	WO 92 06197 A (MINI AGRICULTURE & FISHERIES) 16 April 1992 see claims 2,3 --- -/--	1,2,5,6, 10

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

18 November 1997

Date of mailing of the international search report

02.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Espen, J

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/US 97/12639

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	THORNS CJ ET AL: "The use of latex particle agglutination to specifically detect Salmonella enteritidis." INT J FOOD MICROBIOL, JAN 1994, 21 (1-2) P47-53, NETHERLANDS, XP002047276 see the whole document ---	1,2,5,6, 10
Y	WO 92 06198 A (MINI AGRICULTURE & FISHERIES) 16 April 1992 see claims 1-28 ---	1,2,5,6, 10
Y	WO 93 20231 A (MINI AGRICULTURE & FISHERIES ;WOODWARD MARTIN JOHN (GB); THORNS CH) 14 October 1993 see claims 1-28 ---	1,2,5,6, 10
A	CLOUTHIER SC ET AL: "Characterization of three fimbrial genes, sefABC, of Salmonella enteritidis." J BACTERIOL, MAY 1993, 175 (9) P2523-33, UNITED STATES, XP002047277 -----	

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 97/12639

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9206197 A	16-04-92	AU 660152 B	15-06-95
		AU 8548991 A	28-04-92
		CA 2091982 A	02-04-92
		EP 0551325 A	21-07-93
		JP 6501934 T	03-03-94
		US 5510241 A	23-04-96
		AT 155169 T	15-07-97
		AU 660945 B	13-07-95
		AU 8656691 A	28-04-92
		CA 2091984 A	02-04-92
		DE 69126786 D	14-08-97
		EP 0551324 A	21-07-93
		WO 9206198 A	16-04-92
		JP 6502531 T	24-03-94

WO 9206198 A	16-04-92	AT 155169 T	15-07-97
		AU 660945 B	13-07-95
		AU 8656691 A	28-04-92
		CA 2091984 A	02-04-92
		DE 69126786 D	14-08-97
		EP 0551324 A	21-07-93
		JP 6502531 T	24-03-94
		AU 660152 B	15-06-95
		AU 8548991 A	28-04-92
		CA 2091982 A	02-04-92
		EP 0551325 A	21-07-93
		WO 9206197 A	16-04-92
		JP 6501934 T	03-03-94
		US 5510241 A	23-04-96

WO 9320231 A	14-10-93	AU 3895293 A	08-11-93



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/255, G01N 33/50	A1	(11) International Publication Number: WO 98/03656
		(43) International Publication Date: 29 January 1998 (29.01.98)

(21) International Application Number: PCT/US97/12639

(22) International Filing Date: 18 July 1997 (18.07.97)

(30) Priority Data:
60/022,191 19 July 1996 (19.07.96) US(71) Applicant (for all designated States except US): REGENTS OF
THE UNIVERSITY OF MINNESOTA [US/US]; Morrill
Hall, 100 Church Street, S.E., Minneapolis, MN 55455 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RAJASHEKARA,
Gireesh [IN/US]; Minn. Graduate Club, 2089 Carter Av-
enue, St. Paul, MN 55108 (US). NAGARAJA, Kakambi,
V. [IN/US]; 2235 Milton Street North, Roseville, MN
55113 (US). KAPUR, Vivek [IN/US]; Apartment 108,
2572 Kenzie Terrace, St. Anthony, MN 55418 (US).(74) Agent: BRUESS, Steven, C.; Merchant, Gould, Smith, Edell,
Welter & Schmidt, P.A., 3100 Norwest Center, 90 South
Seventh Street, Minneapolis, MN 55402-4131 (US).(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility
model), DE, DE (Utility model), DK, DK (Utility model),
EE, EE (Utility model), ES, FI, FI (Utility model), GB,
GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility
model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU,
ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB,
GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ,
CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.

(54) Title: RECOMBINANT SEF14 FIMBRIAL PROTEIN FROM SALMONELLA

(57) Abstract

A truncated SE fimbria antigen useful as an antigen for immunoassay diagnosis of *Salmonella enteritidis* (SE) infection or evidence
of infection.

*(Referred to in PCT Gazette No. 27/1998, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

RECOMBINANT SEF14 FIMBRIAL PROTEIN FROM SALMONELLA

Field of the Invention

The present invention relates to a method of
5 cloning and expressing a truncated form of a fimbrial gene
— and the use of the truncated fimbrial gene product in an
immunodiagnostic assay and for immunoprophylaxis.

Background of the Invention

10 Foodborne infections cause an estimated 6.5
million cases of human illness and 9000 deaths annually in
the United States alone. Bacterial infections by
Salmonella are the most commonly reported cause of
foodborne outbreaks. *Salmonella enteritidis* (SE) is the
15 dominant *Salmonella* serotype isolated from cases of food
poisoning. Many of these outbreaks are thought to be due
to infected poultry products, particularly eggs and egg
products.

The best way to prevent infection in human
20 populations is to diagnose and treat the infected animal
prior to human consumption. Because the greatest threat of
food poisoning from *Salmonella* is from poultry products,
there is a need for a method to detect birds that are
infected with SE.

25 Some current diagnostic methods rely on
conventional bacteriologic cultures. However, these
procedures are relatively slow, often taking up to 3 to 4
days to provide even a presumptive diagnosis.
Additionally, the great susceptibility of SE to physical
30 and chemical factors such as desiccation, radiation, low
temperature, heating, or chemical preservatives, causes
traditional bacteriologic culture methods to generally have
a low sensitivity. Consequently, many birds or animals

that are infected with SE are often not detected when conventional bacterial cultures are used.

Other diagnostic methods rely on the detection of serum antibodies specific to SE. Although several serological methods such as micro-agglutination, serum plate agglutination, latex particle agglutination microantiglobulin, ELISA have previously been employed, these assays lack either the sensitivity or specificity necessary to detect SE infected birds, or the tests are too difficult to perform in a routine laboratory or field setting. Consequently, widespread application of these tests for the detection of SE infections has been impractical.

A useful antigenic determinant that is found on many species of *Enterobacteriaceae* are fimbriae, proteinaceous filamentous surface structures composed of protein subunits called fimbrin. Upon infection, birds make antibodies to this SE fimbrial antigen. Therefore, the SE fimbrial antigen is useful in a diagnostic assay for the presence of SE in poultry.

SE is known to have at least four distinct fimbria, designated Sef14, Sef17, Sef18 and Sef21. These proteins are encoded by *SefA*, *AgfA*, *SefD* and *FinA* genes, respectively.

Although the gene encoding Sef14 has been identified and its DNA nucleotide sequence determined (Trucotte and Woodward, *Journal of General Microbiology*, 139:1477-1485 (1993)), an effective diagnostic method using this surface antigen has not been developed, partially due to the difficulty of efficiently producing the fimbriae proteins in purified form and in large quantities. Additionally, expression of Sef14 fimbriae by cultured

Salmonella enteritidis is highly dependent on the growth medium composition. In a study by Thorns et al., *International Journal of Food Microbiology*, 21:47-53 (1994), only peptone water pH 7.2 supported the expression of Sef14 by all *Salmonella enteritidis* strains examined. Consequently, previous diagnostic assays using Sef14 have used antibodies against Sef14 and not the antigen itself.

Hence, there is a need for a sensitive, specific and routine antigen and method to reliably detect SE infection in birds, preferably a method that is easily adaptable to large-scale screening of poultry flocks.

Summary of the Invention

The present invention provides a sensitive, specific, routine antigen and assay to reliably detect SE-infected animals. Specifically, the present invention provides a truncated form of the Sef14 antigen that can be easily produced in purified form and in large quantities and used in the method of the invention. The novel Sef14 antigen, when coupled to a substrate such as latex beads, provides a diagnostic assay for SE, particularly useful in large-scale screening of poultry flocks.

Brief Description of the Figures

Figure 1 is a photograph showing a SDS-PAGE of the recombinant Sef14 (rSef14) fragment (arrow).

Figure 2 is a photograph showing a Western blot of the rSef14 fragment probed with anti-Sef14 antibody (lane 1) and anti-tag (T7) antibody (lane 2).

Figure 3 is a photograph showing results of a rSef14-latex agglutination assay for SE infection in

chickens exposed to *S. enteritidis* (A), *S. pullorum* (B), and serum-free antigen control (C).

Figure 4 is a photograph showing results of a rSef14-latex agglutination assay for SE infection in chickens exposed to *S. enteritidis* (A), *S. gallinarum* (B), *S. pullorum* (C), *S. typhimurium* (D), *C. arizonae* (E), *E. coli* (F), serum free antigen control (G), and serum control (H).

Figure 5 is a graph showing the percentage of chickens testing positive for anti-SE antibodies during 4 weeks post-inneculation. The five bars at each week represent inneculation with 10^4 , 10^6 , 10^8 , 10^{10} , and control (no cells).

Figure 6 is a graph showing the antibody titres of chicken sera samples testing positive for anti-SE antibodies.

Figure 7 is a graph showing the antibody titres of chicken egg yolk samples testing positive for anti-SE antibodies.

Detailed Description of the Invention

The present invention is directed to a method for diagnosing *Salmonella enteritidis* infection or evidence of infection in an animal, particularly poultry, using a recombinant truncated fimbrial antigen.

"Infection" means active colonization of the animal by SE organisms. "Evidence of infection" means a prior history of colonization by SE in the animal, although active colonization is not present. Diagnosis of active infection is needed to protect against contamination of food supplies, whereas diagnosis of prior infection is

needed to alert against new infection or to trace the source of infection in a flock.

Fimbrial Proteins

5 Fimbriae are proteinaceous filamentous surface structures composed of protein subunits called fimbrin. These proteinaceous structures are thought to be virulence factors which mediate specific attachment to host cell mucosal surfaces. They are present in most enteric
10 bacteria capable of invading host cells.

Salmonella enteritidis has four distinct fimbriae: Sef14, Sef17, Sef18 and Sef21 which are encoded by *sefA*, *agfA*, *sefD* and *fimA* genes, respectively. Sef14 is unique with only limited distribution in the genus. In
15 contrast, all other fimbrial proteins are widely distributed in the genus. Thus, they have limited use as diagnostic reagents for SE detection.

Cloning and Expression of Sef14

20 In the present invention, a truncated form of the Sef14 antigen retaining the antigenic character of the entire protein has been produced. Unlike the complete protein, however, the truncated form can be easily produced in purified form and in large quantities, without special
25 growth medium requirements.

PCR technology is used to produce the truncated Sef14 protein by amplification with suitable primers.

Primers are selected to amplify the gene encoding Sef14 in a region downstream of the encoded signal peptide, e.g., downstream of about nucleotide 145 of the *DraI*
30 genomic fragment shown in Figure 1 of Turcotte and Woodward, *Supra*. Preferably, the PCR primers include

additional nucleotides at the 5' ends, encoding specific restriction enzyme recognition sequences, for ease of purification. For example, useful primers for amplifying that portion of the *sefA* gene encoding an immunogenic Sef14 fragment downstream of the signal peptide are shown below:

GGGAATTCGCTGGCTTTGTTGGTAACA	SEQ ID NO:1
GGGCTCGAGTTAGTTTTGATACTGAACGTA	SEQ ID NO:2

After a truncated gene sequence encoding Sef14 is produced, it can be cloned into a host using a plasmid or phage as a vector. Typically, the expression of Sef14 fimbriae by cultured *Salmonella enteritidis* is highly dependent on the growth medium composition (Thorns et al, *International Journal of Food Microbiology*, 21:47-53 (1994)), and it is typically difficult to produce large quantities. However, a truncated form of Sef14 having at least the signal peptide removed is expressed in host systems such as *E. coli* without these difficulties.

Truncated Sef14 Antigen

Because the truncated Sef14 protein retains the antigenic characteristics of the complete protein, it is useful in various immunological methods. For example, the inventive antigen is useful in antibody binding immunoassays such as assays to detect the presence of antibodies against SE in a sample. Suitable binding assays include ELISA, wherein the recombinant Sef14 antigen is bound to a surface and exposed to antibodies against SE. To detect the presence of bound anti-SE antibodies, a marker such as an enzyme-linked secondary antibody is then added.

An agglutination assay using truncated Sef14 antigen-coated latex beads is preferred. In the agglutination reaction, antigen-coated latex beads form detectable clusters when exposed to antibodies against SE. This preferred assay is described more fully in Example 4, below.

Diagnostic Assays

The assays described above can be used to detect the presence of antibodies to *Salmonella enteritidis*. Preferably, the assays are used to determine whether or not an animal, e.g. a poultry animal such as a chicken or turkey, is infected with SE. Animal fluid such as blood or serum can be used in a diagnostic assay. If an animal is infected with SE, the animal will typically produce anti-SE antibodies. The recombinant Sef14 antigen is used to detect the presence of anti-SE antibodies, SE infection or the SE organism itself. Diagnostic assays such as these are particularly useful in birds. More particularly, diagnostic assays are useful in detecting SE infections in chicken or turkey to prevent foodborne illness by poultry consumption.

Vaccine

Passive immunization with anti-Sef14 antibodies has been shown to reduce *Salmonella enteritidis* colonization (Peralta et al. 1994). Additionally, Sef14 can induce a T-cell immune response (Ogunniyi et al 1994). Because the truncated Sef14 antigen exhibits these immunological activities, can be produced in large quantities, and does not have the cumbersome growth requirements of the complete protein, the truncated Sef14

antigen is also useful as a vaccine to confer immunity against SE. Preferably, the truncated Sef14 antigen is used as a vaccine in poultry to prevent foodborne illnesses.

5

EXAMPLES

The invention may be better understood with reference to the following examples which are not intended to limit the invention.

10

Example 1

Isolation of *S. enteritidis* genomic DNA

15

S. enteritidis was grown overnight at 37°C in Luria-Bertani (LB) broth. Genomic DNA was extracted as described (Sambrook, et al., 1989) using standard methods with minor modifications. In brief, bacterial cells were pelleted by centrifugation at 13,000 x g for 3 minutes, washed/suspended in 1 ml of 1 M NaCl, centrifuged for 5 minutes at 13,000 x g, and the pellet resuspended in 1 ml TE buffer (50 mM Tris-HCl, 50 mM EDTA, pH 7.8). The sample was next incubated with 5 µl of lysozyme (50 mg/ml) (Sigma Chemical Co., St. Louis, MO) and 0.3 mg/ml RNase A (Sigma) at 37°C for 30 minutes. To this suspension, 1% sarkosyl and 0.6 mg/ml of proteinase K (Sigma) were added, and the mixture incubated at 37° for 1 hour. Following incubation, chromosomal DNA was extracted twice with an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1) and once with chloroform:isoamyl alcohol (24:1). Genomic DNA in the aqueous phase was precipitated at -20°C with two volumes of absolute ethanol and 0.1 volume of 3 M sodium acetate, and

20

25

30

pelleted by centrifugation at 13,000 x g for 5 minutes. The pellet was then washed twice with 70% ethanol, air dried, and suspended in TE Buffer (10mM Tris-HCl and 1mM EDTA pH 8.0). Total DNA was quantitated spectrophotometrically at A₂₆₀ nm.

Example 2

Cloning of *sefA* gene fragment

Oligonucleotide primer selection and synthesis:

Oligonucleotide primers corresponding to an internal fragment (64-498 bp) of the open-reading frame of the *sefA* gene were used for PCR amplification. Additional bases were added to the 5' end of each primer in order to confer a recognition sequence for either *EcoRI* (forward primer) or *XhoI* (reverse primer). The oligonucleotide primers were obtained from Integrated DNA technologies Inc., Ames, IA. The DNA sequences for the forward and reverse primers are shown below:

<u>GGGAATTC</u> GCTGGCTTTGTTGGTAACA	SEQ ID NO:1
GGGCTCGAGTTAGTTTGTGATACTGAACGTA	SEQ ID NO:2

Additional nucleotides added to the 5' end of the primers are underlined.

PCR amplification of *sefA* gene fragment.

Amplification reactions were performed in 30 µl volumes with 30 pmol of each primer and 5 mM MgCl₂. The reagents and enzymes used for PCR were obtained either from Boehringer Mannheim (Indianapolis, IN) or Perkin Elmer (Foster City, CA). One hundred ng of genomic DNA was used as a template for PCR amplification with the following parameters: an initial denaturation at 94°C for 5 minutes, followed by 35 cycles of denaturation (94°C for 1.5

minutes), annealing 52°C for 1 minute) and extension (72°C for 2 minutes), and a final extension of 15 minutes at 72°C. All amplification reactions were performed in a Perkin-Elmer Cetus DNA thermal cycler (Model 480). The PCR products were analyzed on a 1% agarose gel, stained with ethidium bromide (0.5µg/ml), and photographed under UV light.

PCR products were gel extracted (Qiagen Inc., Chatsworth, CA), quantitated spectrophotometrically, at 260 nm, and cloned directly into pGEM-T vector (Promega, Madison, WI). Following ligation, 2µl of the reaction products were transformed into *E. coli* DH5α cells (Gibco BRL, Gaithersburg, MD) by the heat shock method. Recombinant colonies were selected on ampicillin/IPTG-Xgal containing plates and screened for the presence of the appropriate insert by restriction analysis.

Nucleotide sequence analysis

A bacterial colony containing the recombinant plasmid with the *rSefA* fragment was grown in LB-ampicillin media, and the plasmid extracted using Qiagen plasmid extraction kit (Qiagen). The nucleotide sequence of the insert was determined using oligonucleotide primers specific to the vector sequence by automated DNA sequencing at the University of Minnesota Advanced Genetic Analysis Center. The insert was sequenced in its entirety in both orientations, and the amino acid sequence deduced using the standard genetic code (DNA*, Madison, WI). Sequencing results are shown below for nucleotide and deduced amino acid sequences of the insert (Seq.ID.NO:5), together with a tag sequence added during the subcloning of the fragment into the pET/abc expression vector (Seq.ID.NO:3). The

added tag sequence at the 5' end, provides a Histidine-rich portion to facilitate purification of the sequence on nickel columns, as well as an antigenic region that specifically binds the T7 anti-tag antibody provided with the pET/abc vector kit.

Nucleic Acid Sequence encoding rSefA fragment

SEQ ID NO:3

ATG GGC AGC AGC CAT CAT CAT CAT CAT CAC AGC AGC GGC CTG GTG	45
CCG CGC GGC AGC CAT ATG GCT AGC ATG ACT GGT GGA CAG CAA ATG	90
GGT CGC GGA TGG GAA TTC GCT GGC TTT GTT GGT AAC AAA GCA GTG	135
GTT CAG GCA GCG GTT ACT ATT GCA GCT CAG AAT ACA ACA TCA GCC	180
AAC TGG AGT CAG GAT CCT GGC TTT ACA GGG CCT GCT GTT GCT GCT	225
GGT CAG AAA GTT GGT ACT CTC AGC ATT ACT GCT ACT GGT CCA CAT	270
AAC TCA GTA TCT ATT GCA GGT AAA GGG GCT TCG GTA TCT GGT GGT	315
GTA GCC ACT GTC CCG TTC GTT GAT GGA CAA GGA CAG CCT GTT TTC	360
CGT GGG CGT ATT CAG GGA GCC AAT ATT AAT GAC CAA GCA AAT ACT	405
GGA ATT GAC GGG CTT GCA GGT TGG CGA GTT GCC AGC TCT CAA GAA	450
ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT AAA TCG ACC CTG CCA	495
GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG CAG TAT CAA AAC	540
TAA <u>CTC GAG CCC</u> 552	

*Additional amino acid residues added to the amino terminus to facilitate protein purification and cloning are underlined.

Deduced amino acid sequence of rSefA protein fragment*

Seq. ID NO: 4

MGSSHHHHHHSSGLVPRGSHMASMTGGQOMGRGSEFAGFVG NKAVVQAAVT
IAAQNTTSANWSQDPGFTGPAVAAGQKVGTLSITATGPHNSVSIAGKGASVSGG
VATVPFVDGQGQPVFRGRIQGANINDQANTGIDGLAGWRVASSQETLNVPVTT
FGKSTLPAGTFTATFYVQQYQN

*Additional amino acid residues added to the amino terminus to facilitate protein purification and cloning are underlined.

It is understood that the amino acids added to the N-terminus of the Spf14 antigen are optional, and used for ease of cloning and purification. The amino acid sequence in the absence of these added residues (Sequence ID No:6)

with or without other added residues for cloning or purification procedures, for example, are similarly useful as antigens in the diagnostic assays of the invention.

5 **Subcloning *sefA* gene fragment into an expression vector**

 The pGEM-T plasmid carrying *sefA* fragment was double digested with *EcoRI* and *XhoI*, and the digested products gel purified (Qiagen) and cloned into *EcoRI* and *XhoI* digested pET/abc expression vectors (Novagen Inc.,
10 Madison, WI). Ligation products (2 μ l) from each of the reactions were transformed into *E. coli* BL21(DE3)pLyS cells by heat shock method. Recombinant clones were cultured on kanamycin and chloramphenicol containing plates, and analyzed by restriction enzyme analysis.

15

rSefA fragment expression

 The recombinant clones were selected based on restriction enzyme analysis with *EcoRI* and *XhoI* digestion, selecting those clones yielding appropriately sized
20 fragments as compared with a vector control. Selected clones were analyzed for rSefA fragment expression. Briefly, a single colony from each (pETabc/SefA fragment) freshly streaked plate was picked and inoculated to 50 ml LB broth containing appropriate antibiotics and incubated
25 with shaking at 200 rpm at 37°C until the OD₆₀₀ reached 0.6. Cultures were induced with IPTG (0.4 mM) and incubated for an additional 3 hours. Following incubation, the cells were pelleted and resuspended in 5 ml of TE buffer (50mM
30 Tris-HCl pH 8.0, 2mM EDTA) and incubated with 25 μ l of lysozyme (50 mg/ml) and 100 μ l of 1% Triton X-100 for 20 minutes at 30°C. The samples were sonicated until they were no longer viscous, and centrifuged at 39,000 x g for

20 minutes. The supernatant was passed through a 0.45 μ m membrane filter, and stored at -20°C until further use.

SDS-PAGE analysis

5 The cell lysates were next analyzed by SDS-PAGE for the presence of the rSefA fragment by mixing with an equal volume of 2x SDS solubilization buffer separating on 12% polyacrylamide gels, and staining with Coomassie blue. The results are shown in lane 1 of Figure 1 which contains
10 the total protein produced by the vector and contained in the cell lysates.

Western blot analysis

 The lysates were separated on 12% polyacrylamide
15 gels and transferred onto a nitrocellulose membrane using Transblot apparatus (Bio-Rad laboratories, Hercules, CA). Following transfer, the membrane was blocked with 3% BSA in phosphate buffered saline (PBS) and stained with either T7 anti-tag antibody (Novagen) or rabbit anti-Sef14 specific
20 antibody (kindly provided by Dr. W. W. Kay, University of Victoria, BC, Canada). The membrane was washed and stained with anti-rabbit IgG/HRP conjugate and treated with
developing reagent (Amersham lif sciences, Inc., USA) for 1 minute, exposed to X-ray film, and the radiograph
25 developed. The results are shown in Figure 2, where lane 1 is probed with anti-Sef14 antibody, and lane 2 with T7 anti-tag antibody.

Purification of rSef14 fragment protein by column 30 chromatography and electroelution

 The recombinant Sef14 protein fragment produced in the cell lysates described above was purified by binding

of the Histidine-rich tag to nickel columns as described by the manufacturer (Novagen). Briefly, the cells were induced and extract was prepared as described above except that the induced cells were suspended in Tris buffer without EDTA. The cell lysate was passed through nickel columns and washed sequentially with binding buffer (5 mM imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9) and wash buffer (60 mM imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9). The bound protein was eluted using elution buffer (1 M imidazole, 0.5 M NaCl, 20 mM Tris-HCl, pH 7.9), quantitated using a Bio-Rad protein assay kit (Bio-Rad laboratories, Hercules, CA), and analyzed by SDS-PAGE. (See Figure 1, lane 2, where the arrow indicates the rSef14 fragment at about 19 KDa.) Since the column purified recombinant material contained traces of non-specific proteins, the appropriate rSef14 fragment was further purified by cutting the rSef14 fragment from the gel and electroelution (Bio-Rad) following the manufacturer's suggested protocol. The electroeluted fragment is shown in lane 4 of Figure 1 (at arrow).

Example 3

Covalent coupling of rSef14 to blue-dyed latex beads

The electroeluted rSef14 protein fragment was coupled to either 0.5 μ m or 1.0 μ m blue-dyed latex beads (Polysciences Inc., Warrington, PA) by gluteraldehyde method. Briefly, 1 ml of 2.5% suspension of the beads were washed with PBS (pH 7.4), pelleted by centrifugation and resuspended in 1 ml of 8% gluteraldehyde (EM grade) in PBS, and incubated overnight with gentle end-to-end mixing at room temperature. Following gluteraldehyde treatment, the beads were pelleted, washed with PBS three times and

incubated with 500 µg of purified rSef14 fragment for 5 hours at room temperature with gentle end-to-end mixing. The beads were pelleted, and incubated with 1 ml of 0.5 M ethanolamine in PBS for 30 minutes at room temperature with gentle end-to-end mixing. The mixture was then treated with 1 ml of 10 mg/ml BSA in PBS for 30 minutes at room temperature, centrifuged and the pellet resuspended in 1 ml PBS (pH 7.4), containing 10 mg/ml BSA, 0.1% NaN₃, and 5% glycerol, and stored at 4°C to form rSef14 - fragment coated latex beads for use in agglutination assays.

Example 4

rSef14-latex bead agglutination test

Bacteria was administered to chickens by either injection, intratracheal or oral administration of 10⁷ colony forming units (CFU) of either *S. enteritidis*, *S. pullorum*, *S. arizonae*, *S. typhimurium*, *S. gallinarum*, or *E. coli*. After about two to three weeks exposure, serum was collected and used to evaluate the sensitivity and specificity of the rSef14-latex beads in an agglutination assay for anti-SE antibody binding. A total volume of 7.5 µl of rSef14 fragment coated latex beads, produced as described for Example 3, were mixed with an equal volume of chicken serum collected from birds exposed to various pathogens, as described above. The presence of agglutination, visually seen as a loss of intense blue color in the sample (i.e., lightening of color as the coated beads agglutinate or form a lattice). Absence of the agglutination reaction was visualized by the remaining intense blue color of the dyed beads in a homogeneous suspension. Positive or negative agglutination reaction

was recorded after two minutes. The results are shown in Figures 3 and 4.

In figure 3, intense blue color (negative result) is seen in test samples B and C (*S. pullorum* and the serum-free antigen control). In contrast, a positive agglutination result is seen in test Sample A, (*S. enteritidis*), as a pale blue, diffuse agglutination pattern.

In figure 4, a positive agglutination reaction is seen in sample A (*S. enteritidis*) and in sample H (serum control). No agglutination reaction is seen in the samples B-G containing serum animals exposed to the following pathogens: *S. gallinarium* (B), *S. pullorum* (C), *S. typhimurium* (D), *S. arizonae* (E), *E. coli* (F), and serum free antigen control (H).

Example 5

Detection of anti-S.E. antibodies in infected chickens

To confirm the specificity of the assay of the invention, forty SPF chickens (age 4weeks) were inoculated with various species of *Salmonella*. A suspension of 10^9 CFU in PBS was administered by injection. A booster dose of 10^9 CFU was administered orally two weeks later. Serum samples were taken at weekly intervals and assayed for the presence of anti-SE antibodies.

Serum samples were assayed by the standard Serum Plate Test (SPT) as described in Schaffer and MacDonald, 1931, _____, ____:236-240. Slides were visually scored (+ or -) for the presence of agglutination, indicating antigen-antibody binding. Samples were also assayed by the Microtiter Plate Test (MPT) as described in Williams and Whittemore, 1971 *Applied Microbiology* 21:394-399. These

two assays are standard screening methods for the detection of *Salmonella*, using *S.pullorium* as a whole-cell antigen, and are not specific for SE, as shown in the table below.

To demonstrate the specificity of the assays of the invention, serum samples were assayed using the latex agglutination test (LAT) described above for Example 4, which utilized the truncated Spf14 antigen coupled to latex beads. Serum samples were also assayed for anti-SE antibodies by ELISA. In the ELISA, the truncated Spf14 antigen prepared as described for Example 3, was coated onto polystyrene plates. Antigen-coated plates were exposed to serum samples to permit binding of anti-SE antibodies to the antigen. The bound antigen-antibody complexes were washed, and then incubated with anti-chicken antibody coupled to biotin. The complex was then exposed to strep-avidin for signal detection.

Results are shown in the table below. The LAT and Elisa assays demonstrated a useful specificity for the detection of SE. Of the organisms tested, only *S.dublin*, a bovine pathogen, demonstrated cross-reactivity in the assays.

Species	SPT	MT	LAT	ELISA
<i>S. enteritidis</i>	+	+	+	+
<i>S. gallinarum</i>	+	+	-	-
<i>S. pullorum</i>	+	+	-	-
<i>S. dublin</i>	+	+	+	+
<i>S. berta</i>	+	+	-	-
<i>S. typhimurium</i>	-	+	-	-
<i>E. coli</i>	-	-	-	-
Control (no cells)	-	-	-	-

Example 6

Specificity of anti-SE assay

The ELISA assay for detecting anti-SE antibodies described above for Example 6 was tested for specificity using a panel of antisera against known pathogenic organisms. Each sera was assayed in the anti-SE ELISA. No crossreactivity was observed with any of the tested antisera.

Antisera	ELISA		Antisera	ELISA
Pox	-		MG	-
Reo	-		NDV	-
Rev	-		CAV	-
SB-1	-		HVT	-
IBDV	-		IBV	-
ILT	-		<i>S.typhimurum</i>	-
LLA	-		<i>S.gallinarum</i>	-
LLB	-		<i>S.pullorum</i>	-
MS	-			

Example 7

Sensitivity of ELISA for detection of SE

Fifty white leghorn layer chickens (5 weeks old) were orally inoculated in a single exposure with varied amounts of SE, from 10^4 to 10^{10} CFU in PBS. Serum samples were collected at weekly intervals for up to seven weeks. Eggs were collected for egg yolk antibody detection.

Samples were analyzed for detection of anti-SE

antibodies using the ELISA described above for Example 6. As shown in Figure 5, control chickens showed no positive reaction in the ELISA assay. Approximately 40-80% of chickens exposed to 10^4 , 10^6 , 10^8 , and 10^{10} CFU of SE tested positive for anti-SE antibodies during the first four weeks post-inneculation. From 4-7 weeks post-inneculation, the data stabilized at about 45% positive detection of anti-SE antibodies.

Antibody titers in the sera and egg yolks of chickens exposed to 10^4 , 10^6 , and 10^8 CFU of SE and testing positive in the ELISA for anti-SE antibodies are shown in Figures 6 and 7.

These data demonstrate specific detection of anti-SE antibodies using recombinant Sef14-antigen coated latex beads in an agglutination assay and using the antigen as a capture agent in an ELISA. These assays provide a sensitive and specific diagnostic tool for the detection of anti-SE antibodies in animals and for the diagnosis of SE infection.

20

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: REGENTS OF THE UNIVERSITY OF MINNESOTA
- (ii) TITLE OF THE INVENTION:
RECOMBINANT FIMBRIAL PROTEIN
- (iii) NUMBER OF SEQUENCES: 6
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Merchant, Gould, Smith, Edell, Welter & Schmidt
 - (B) STREET: 3100 Norwest Center, 90 South Seventh St
 - (C) CITY: Minneapolis
 - (D) STATE: MN
 - (E) COUNTRY: USA
 - (F) ZIP: 55402
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 18-JUL-1997
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/022,191
 - (B) FILING DATE: 19-JUL-1996
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Kettelberger, Denise M
 - (B) REGISTRATION NUMBER: 33,924
 - (C) REFERENCE/DOCKET NUMBER: 600.335W001
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 612/371-5268
 - (B) TELEFAX: 612/332-9081
 - (C) TELEX:

21

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGGAATTCGC TGGCTTTGTT GGTAACA

27

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GGGCTCGAGT TAGTTTTGAT ACTGAACGTA

30

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 552 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 1...540
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATG GGC AGC AGC CAT CAT CAT CAT CAC AGC AGC GGC CTG GTG CCG 48
 Met Gly Ser Ser His His His His His His Ser Ser Gly Leu Val Pro
 1 5 10 15

CGC GGC AGC CAT ATG GCT AGC ATG ACT GGT GGA CAG CAA ATG GGT CGC 96
 Arg Gly Ser His Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg
 20 25 30

22

GGA TGG GAA TTC GCT GGC TTT GTT GGT AAC AAA GCA GTG GTT CAG GCA Gly Trp Glu Phe Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala 35 40 45	144
GCG GTT ACT ATT GCA GCT CAG AAT ACA ACA TCA GCC AAC TGG AGT CAG Ala Val Thr Ile Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln 50 55 60	192
GAT CCT GGC TTT ACA GGG CCT GCT GTT GCT GCT GGT CAG AAA GTT GGT Asp Pro Gly Phe Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly 65 70 75 80	240
ACT CTC AGC ATT ACT GCT ACT GGT CCA CAT AAC TCA GTA TCT ATT GCA Thr Leu Ser Ile Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala 85 90 95	288
GGT AAA GGG GCT TCG GTA TCT GGT GGT GTA GCC ACT GTC CCG TTC GTT Gly Lys Gly Ala Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val 100 105 110	336
GAT GGA CAA GGA CAG CCT GTT TTC CGT GGG CGT ATT CAG GGA GCC AAT Asp Gly Gln Gly Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn 115 120 125	384
ATT AAT GAC CAA GCA AAT ACT GGA ATT GAC GGG CTT GCA GGT TGG CGA Ile Asn Asp Gln Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg 130 135 140	432
GTT GCC AGC TCT CAA GAA ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT Val Ala Ser Ser Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly 145 150 155 160	480
AAA TCG ACC CTG CCA GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG Lys Ser Thr Leu Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln 165 170 175	528
CAG TAT CAA AAC TAACTCGAGC CC Gln Tyr Gln Asn 180	552

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 180 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

23

```

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
 1          5          10          15
Arg Gly Ser His Met Ala Ser Met Thr Gly Gly Gln Gln Met Gly Arg
          20          25          30
Gly Trp Glu Phe Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala
 35          40          45
Ala Val Thr Ile Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln
 50          55          60
Asp Pro Gly Phe Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly
65          70          75          80
Thr Leu Ser Ile Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala
          85          90          95
Gly Lys Gly Ala Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val
          100          105          110
Asp Gly Gln Gly Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn
          115          120          125
Ile Asn Asp Gln Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg
          130          135          140
Val Ala Ser Ser Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly
145          150          155          160
Lys Ser Thr Leu Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln
          165          170          175
Gln Tyr Gln Asn
          180

```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 435 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(ix) FEATURE:

- (A) NAME/KEY: Coding Sequence
- (B) LOCATION: 1...432
- (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

GCT GGC TTT GTT GGT AAC AAA GCA GTG GTT CAG GCA GCG GTT ACT ATT      48
Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala Ala Val Thr Ile
 1          5          10          15

GCA GCT CAG AAT ACA ACA TCA GCC AAC TGG AGT CAG GAT CCT GGC TTT      96
Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln Asp Pro Gly Phe
          20          25          30

ACA GGG CCT GCT GTT GCT GCT GGT CAG AAA GTT GGT ACT CTC AGC ATT      144
Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly Thr Leu Ser Ile
          35          40          45

```

24

ACT GCT ACT GGT CCA CAT AAC TCA GTA TCT ATT GCA GGT AAA GGG GCT 192
 Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala Gly Lys Gly Ala
 50 55 60

TCG GTA TCT GGT GGT GTA GCC ACT GTC CCG TTC GTT GAT GGA CAA GGA 240
 Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val Asp Gly Gln Gly
 65 70 75 80

CAG CCT GTT TTC CGT GGG CGT ATT CAG GGA GCC AAT ATT AAT GAC CAA 288
 Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn Ile Asn Asp Gln
 85 90 95

GCA AAT ACT GGA ATT GAC GGG CTT GCA GGT TGG CGA GTT GCC AGC TCT 336
 Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg Val Ala Ser Ser
 100 105 110

CAA GAA ACG CTA AAT GTC CCT GTC ACA ACC TTT GGT AAA TCG ACC CTG 384
 Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly Lys Ser Thr Leu
 115 120 125

CCA GCA GGT ACT TTC ACT GCG ACC TTC TAC GTT CAG CAG TAT CAA AAC T 433
 Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln Gln Tyr Gln Asn
 130 135 140

AA 435

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ala Gly Phe Val Gly Asn Lys Ala Val Val Gln Ala Ala Val Thr Ile
 1 5 10 15
 Ala Ala Gln Asn Thr Thr Ser Ala Asn Trp Ser Gln Asp Pro Gly Phe
 20 25 30
 Thr Gly Pro Ala Val Ala Ala Gly Gln Lys Val Gly Thr Leu Ser Ile
 35 40 45

Thr Ala Thr Gly Pro His Asn Ser Val Ser Ile Ala Gly Lys Gly Ala
 50 55 60
 Ser Val Ser Gly Gly Val Ala Thr Val Pro Phe Val Asp Gly Gln Gly
 65 70 75 80
 Gln Pro Val Phe Arg Gly Arg Ile Gln Gly Ala Asn Ile Asn Asp Gln
 85 90 95
 Ala Asn Thr Gly Ile Asp Gly Leu Ala Gly Trp Arg Val Ala Ser Ser
 100 105 110
 Gln Glu Thr Leu Asn Val Pro Val Thr Thr Phe Gly Lys Ser Thr Leu
 115 120 125
 Pro Ala Gly Thr Phe Thr Ala Thr Phe Tyr Val Gln Gln Tyr Gln Asn
 130 135 140

WE CLAIM:

1. A method for detecting anti- *Salmonella enteritidis* antibodies in animals, the method comprising:

5 reacting a sample obtained from an animal with a truncated Sef14 antigen under conditions to permit anti-Sef14 antibodies to bind the antigen, the truncated antigen having at least the native Sef14 signal peptide removed; and

10 correlating antibody-antigen binding with the presence of anti-SE antibodies in the sample.

2. A method for diagnosing *Salmonella enteritidis* infection in animals, the method comprising:

15 reacting a sample obtained from an animal with a truncated Sef14 antigen under conditions to permit anti-Sef14 antibodies to bind the antigen, the truncated antigen having at least the native Sef14 signal peptide removed; and

20 correlating antibody-antigen binding with *Salmonella enteritidis* infection.

3. The method of claim 1, wherein the truncated Sef14 antigen has the amino acid sequence of Sequence ID. No. 4.

25 4. The method of claim 1, wherein the truncated Sef14 antigen has the amino acid sequence of Sequence ID. No.6.

5. The method of claim 1, wherein said antigen is fixed to
30 an inert surface prior to said reacting.

6. A Sef14 antigen consisting essentially of the amino acid sequence of Sequence I.D. No. 6.

7. An assay kit for the detection of anti-*Salmonella enteritidis* antibodies comprising an Sef14 antigen consisting essentially of the amino acid sequence of Sequence ID No. 6.

8. The assay kit of claim 7, wherein the antigen consists essentially of the amino acid sequence of Sequence ID.. No. 4.

9. An antigen for stimulating the production of anti-*Salmonella enteritidis* antibodies comprising the amino acid sequence of Sequence ID No. 4 or 6.

10. The method of detecting anti-*Salmonella enteritidis* antibodies described in any of the foregoing claims, wherein the animal samples are obtained from fowl, and particularly from chickens or turkeys.

1/7

FIG. 1


SDS-PAGE of rSef14 fragment protein

FIG. 2

**Western blot - rSef14 probed with Sef14
monospecific polyclonal antibodies**




FIG. 3


Latex agglutination test



(A) (B) (C)


FIG. 4

Latex agglutination test




5/7

FIG. 5



6/7


FIG. 6



SUBSTITUTE SHEET (RULE 26)

7/7

FIG. 7



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/12639

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/31 C07K14/255 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THORNS CJ ET AL: "Development and application of enzyme-linked immunosorbent assay for specific detection of Salmonella enteritidis infections in chickens based on antibodies to SEF14 fimbrial antigen." J CLIN MICROBIOL, APR 1996, 34 (4) P792-7, UNITED STATES, XP002047275 see page 792 - page 793; table 1 ---	1,2,5,6, 10
X	WO 92 06197 A (MINI AGRICULTURE & FISHERIES) 16 April 1992 see claims 2,3 --- -/--	1,2,5,6, 10

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

18 November 1997

Date of mailing of the international search report

02.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Espen, J

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/US 97/12639

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	THORNS CJ ET AL: "The use of latex particle agglutination to specifically detect Salmonella enteritidis." INT J FOOD MICROBIOL, JAN 1994, 21 (1-2) P47-53, NETHERLANDS, XP002047276 see the whole document ---	1,2,5,6, 10
Y	WO 92 06198 A (MINI AGRICULTURE & FISHERIES) 16 April 1992 see claims 1-28 ---	1,2,5,6, 10
Y	WO 93 20231 A (MINI AGRICULTURE & FISHERIES ;WOODWARD MARTIN JOHN (GB); THORNS CH) 14 October 1993 see claims 1-28 ---	1,2,5,6, 10
A	CLOUTHIER SC ET AL: "Characterization of three fimbrial genes, sefABC, of Salmonella enteritidis." J BACTERIOL, MAY 1993, 175 (9) P2523-33, UNITED STATES, XP002047277 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/12639

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9206197 A	16-04-92	AU 660152 B	15-06-95
		AU 8548991 A	28-04-92
		CA 2091982 A	02-04-92
		EP 0551325 A	21-07-93
		JP 6501934 T	03-03-94
		US 5510241 A	23-04-96
		AT 155169 T	15-07-97
		AU 660945 B	13-07-95
		AU 8656691 A	28-04-92
		CA 2091984 A	02-04-92
		DE 69126786 D	14-08-97
		EP 0551324 A	21-07-93
		WO 9206198 A	16-04-92
		JP 6502531 T	24-03-94
WO 9206198 A	16-04-92	AT 155169 T	15-07-97
		AU 660945 B	13-07-95
		AU 8656691 A	28-04-92
		CA 2091984 A	02-04-92
		DE 69126786 D	14-08-97
		EP 0551324 A	21-07-93
		JP 6502531 T	24-03-94
		AU 660152 B	15-06-95
		AU 8548991 A	28-04-92
		CA 2091982 A	02-04-92
		EP 0551325 A	21-07-93
		WO 9206197 A	16-04-92
		JP 6501934 T	03-03-94
		US 5510241 A	23-04-96
WO 9320231 A	14-10-93	AU 3895293 A	08-11-93